

CAUTION

**KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

FARMALINX

Snail Trail

SNAIL AND SLUG PELLETS

ACTIVE CONSTITUENT: 15 g/kg METALDEHYDE

For the control of slugs and snails in agricultural and horticultural crops, vineyards and other agricultural, industrial & commercial situations where snails and slugs congregate.

DIRECTIONS FOR USE:

Restraint: DO NOT heap pellets.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Agricultural Crops Cereals, oilseeds (including poppies) Horticultural Crops Fruit (including grapevines), vegetables	Snails and Slugs including: Common white snail (<i>Cermea virgata</i>) White Italian snail (<i>Theba pisana</i>) Conical snail (<i>Cochlicella acuta</i>) Grey field slug (<i>Deroceras reticulatum</i>) Brown field slug (<i>Deroceras invadens</i>)	10 kg/ha	Broadcast pellets evenly over the surface of the ground where snails and slugs are active. Brown field slug mortality may be delayed, requiring longer feeding times.
Pulses, Pasture establishment	(excludes small brown snails)		Snail Trail is to be used only during the establishment stages of pulse crops and pastures. Broadcast pellets evenly over the crop area to be protected either prior to planting, post-plant pre-emergence or early post emergence (prior to 4-leaf stage)
Other agricultural, industrial & commercial situations where snails and slugs congregate			Broadcast pellets evenly over the surface of the ground where snails and slugs are active.
Around crops and/or border areas where slugs and snails congregate	Slugs, Snails	50 g/m ²	Scatter the pellets around the area to be protected and/or where slugs and snails congregate. Spread pellets evenly.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST

FRUIT AND VEGETABLES: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.

PULSES, OILSEEDS, WINTER CEREALS: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING

PASTURES, PULSES, OILSEEDS, WINTER CEREALS: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 5 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Scatter pellets using an appropriate applicator. Where attack is heavy it may be necessary to repeat the application. The best time to apply is immediately after rain or during humid overcast weather when slugs and snails are most active. The pellets eventually decompose leaving no harmful residue. Use regularly to control reinfestation.

TIMING

The best time to apply is immediately after rain, or during humid overcast weather when slugs and snails are most active. For broadacre crops, start when rain/moisture triggers snail activity in Autumn and before significant egg laying.

PRECAUTION

DO NOT apply to edible plant parts. DO NOT graze or cut for stockfeed any crops while Snail Trail pellets are visible on the plant foliage.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF DOMESTIC ANIMALS AND LIVESTOCK

Keep away from animals, poultry, ducks and domestic pets, as accidental poisoning may be fatal. Dogs and other animals find this bait attractive and it may kill them. If pets or livestock are poisoned contact a veterinary surgeon.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers.

Shake and empty contents into applicator equipment. DO NOT dispose of unused product on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Snail Trail Snail and Slug Pellets is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

APVMA Approval No.: 65053/103674

farmalinx *Partners in Agriculture*

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844 W: www.farmalinx.com



Batch No.

Date of Manufacture

NET CONTENTS
500 kg